

A shallow though extensive H₂ 2.122 μ m imaging survey of Taurus-Auriga-Perseus: I. NGC1333, L1455, L1448 and B1

C.J. Davis¹, P. Scholz^{1,2}, P. Lucas³, M.D. Smith⁴ and A. Adamson¹

¹*Joint Astronomy Centre, 660 North A'ohōkū Place, University Park, Hilo, Hawaii 96720, USA*

²*Department of Physics and Astronomy, University of Victoria, Victoria BC, Canada*

³*Centre for Astrophysics Research, Science & Technology Research Institute, University of Hertfordshire, Hatfield AL10 9AB, UK*

⁴*Centre for Astrophysics & Planetary Science, School of Physical Sciences, The University of Kent, Canterbury CT2 7NR, England*

Accepted 2007 ... ; Received 2007 ... ; in original form 2007 ...

ABSTRACT

We discuss wide-field near-IR imaging of the NGC1333, L1448, L1455 and B1 star forming regions in Perseus. The observations have been extracted from a much larger narrow-band imaging survey of the Taurus-Auriga-Perseus complex. These H₂ 2.122 μ m observations are complemented by broad-band K imaging, mid-IR imaging and photometry from the Spitzer Space Telescope, and published submillimetre CO J=3-2 maps of high-velocity molecular outflows. We detect and label 85 H₂ features and associate these with 26 molecular outflows. Three are parsec-scale flows, with a mean flow lobe length exceeding 11.5'. 37 (44%) of the detected H₂ features are associated with a known Herbig-Haro object, while 72 (46%) of catalogued HH objects are detected in H₂ emission. Embedded Spitzer sources are identified for all but two of the 26 molecular outflows. These candidate outflow sources all have high near-to-mid-IR spectral indices (mean value of $\alpha \sim 1.4$) as well as red IRAC 3.6 μ m-4.5 μ m and IRAC/MIPS 4.5 μ m-24.0 μ m colours: 80% have [3.6]-[4.5] > 1.0 and [4.5]-[24] > 1.5. These criteria – high α and red [4.5]-[24] and [3.6]-[4.5] colours – are powerful discriminants when searching for molecular outflow sources. However, we find no correlation between α and flow length or opening angle, and the outflows appear randomly orientated in each region. The more massive clouds are associated with a greater number of outflows, which suggests that the star formation efficiency is roughly the same in each region.

Key words: stars: formation – infrared: stars – ISM: jets and outflows – ISM: kinematics and dynamics – ISM: individual: Perseus

1 INTRODUCTION

The U.K. Infrared Deep Sky Survey (UKIDSS) is a legacy survey being conducted at the United Kingdom Infrared Telescope (Lawrence et al. 2007). UKIDSS comprises five surveys, ranging from very deep, narrow studies of extragalactic fields, to shallow, broad surveys at both low and high galactic latitudes. One such survey, the Galactic Plane Survey (GPS), will by 2012 observe 1868 deg² of the northern and equatorial galactic plane between $-5^\circ < b < +5^\circ$, in J, H and K down to limiting magnitudes of 20.0, 19.1 and 19.0, respectively (Lucas et al. 2008). The GPS also includes a component covering the Taurus-Auriga-Perseus (T-A-P) complex of star-forming clouds, through narrow-band H₂ 1-0S(1) and broad-band K filters. To date, 140 deg² of the T-A-P have been observed as part of the GPS. Observations of an additional 40 deg² are currently being sought to complete the proposed survey area, which is shown in

Fig. 1. The existing data are available from the WFCAM science archive (see Lucas et al. 2008, for details of how to access these data) and, as mosaicked images, from a dedicated database hosted by the Joint Astronomy Centre¹ using an easy-to-navigate clickable map. The use of the latter is recommended for those wishing to quickly access imaging data to search for H₂ emission features.

The goal of this paper is to present demonstration science from the T-A-P H₂ survey. We therefore focus on just the western end of Perseus, comparing the narrow-band imaging data with photometry from the Spitzer Space Telescope legacy programme *From Molecular Cores to Planet-Forming Disks*, also known as *c2d* (Evans et al. 2003). An area covering almost 4 deg² (representing five WFCAM

¹ <http://www.jach.hawaii.edu/UKIRT/TAP>




Figure 1. An overview of the Taurus-Auriga-Perseus complex. The grey-scale shows integrated CO J=1-0 emission from a survey of the Milky Way obtained with 1.2 m telescopes in Cambridge, MA and on Cerro Tololo, Chile (Dame, Hartmann & Thaddeus 2001). The contours represent visual extinction, A_v , constructed from optical star-counts using Digitised Sky Survey data (Dobashi et al. 2005). The black lines outline the complete region to be observed in H₂ 2.122 μ m emission and broad-band K.

tiles) has been searched for H₂ 2.122 μ m line emission features. The mid-IR data are then used to identify candidate protostars for the molecular outflows found in H₂. We also search for H₂ counterparts of the known Herbig-Haro (HH) objects catalogued by Walawender, Bally & Reipurth (2005a), and compare our images with the CO outflow maps of Knee & Sandell (2000) and Hatchell, Fuller & Richer (2007a). Our aim is to establish the relative effectiveness of optical, near-IR and CO surveys of outflows.

The Perseus cloud complex spans more than 30 pc at a distance of between 200-350 pc (de Zeeuw et al. 1999; Černis 1990). The region was first mapped in CO by Sargent (1979); molecular line and extinction maps have since been presented by Ridge et al. (2006). The early search for Young Stellar Objects (YSOs) and protostars from IRAS data by Ladd, Lada & Myers (1993) has recently been superseded by the Spitzer surveys of Jørgensen et al. (2006), Lada et al. (2006) and Gutermuth et al. (2008). Perseus harbours two particularly active star forming regions, NGC1333 – centred on the bright and very well studied HH complex HH 7-11 (Hodapp & Ladd 1995; Bally et al. 1996; Chrysostomou et al. 2000) – and IC 348, a rich, embedded stellar cluster (Luhman et al. 1998, 2005; Lada et al. 2006). Here we focus only on NGC1333 and surrounding clouds – a region we refer to as “Perseus-West” – although the full extent of Perseus will be imaged in H₂ as part of the T-A-P survey (Fig. 1), and in JHK by the UKIDSS Galactic Clusters Survey (Lawrence et al. 2007).




Figure 2. Colour-magnitude diagram of c2d Perseus data. Black dots represent the entire c2d Perseus catalogue; red circles mark those identified as YSOs, while blue crosses mark those labelled “red” in the catalogue.

2 OBSERVATIONS

2.0.1 WFCAM data

The near-IR wide-field camera WFCAM (Casali et al. 2007) contains four Rockwell Hawaii-II (HgCdTe 2048x2048) arrays spaced by 94% in the focal plane. The pixel scale measures 0.40". To observe a contiguous square field on the sky covering 0.75 deg², a region known as a WFCAM “tile”, observations at four positions are required. At each of these four positions eight exposures are obtained in each filter, with a 20 sec exposure time in H₂ and a 5 sec exposure time in K. The images are shifted by an integral multiple of 0.2" so that in the final, mosaicked images the pixel scale is 0.2"; in this way the sub-arcsecond seeing typically obtained at Mauna Kea is fully sampled and bad pixels may be compensated for.

All WFCAM data are initially reduced by the Cambridge Astronomical Survey Unit (CASU) and are distributed through a dedicated archive hosted by the Wide Field Astronomy Unit (WFAU). The CASU reduction steps are described in detail by Dye et al. (2006) and Irwin et al. (2008); astrometric and photometric calibrations are achieved using 2MASS (Dye et al. 2006; Hewett et al. 2006). Data are then downloaded in bulk from WFAU and combined into tiles using astrometric information stored in the file headers. As part of this process, residual sky/background structure is removed by fitting a coarse surface to each image (as described by Davis et al. 2007). Currently data for 140 deg² are available; H₂ and K-band mosaic tiles are available to all interested parties from a database at the Joint Astronomy Centre (<http://jach.hawaii.edu/UKIRT/TAP>) as Rice-compressed fits files.

2.0.2 Spitzer data

The Spitzer Space Telescope observations discussed in this paper were obtained with the IRAC and MIPS cameras, which are described by Fazio et al. (2004) and Rieke et al. (2004), respectively. IRAC data in bands 1,2,3 and 4 (at 3.6, 4.5, 5.8 and 8.0 μm) and MIPS band 1 data (at 24 μm) were obtained in September 2004 as part of the Spitzer legacy programme *c2d* (Evans et al. 2003). The data have been presented by Jørgensen et al. (2006); guaranteed time observations of NGC1333 are presented by Gutermuth et al. (2008). Additional data on L1448 were obtained on February 25, 2005 by Tobin et al. (2007).

The mid-IR photometry used here were obtained directly from the *c2d* archive (<http://peggysue.as.utexas.edu/SIRTF/>), which includes data extracted from the third delivery of IRAC and MIPS data. The procedures used during source extraction are described by Evans et al. (2005) and Harvey et al. (2007). For Perseus, the full catalogue of sources detected in at least one IRAC band contains over 120,000 objects. IRAC and MIPS band 1 (24 μm) data have been combined with 2MASS JHK photometry by the *c2d* group to distinguish candidate Young Stellar Objects (YSOs) and protostars from field stars and background galaxies (see also Jørgensen et al. 2006). A spectral index, $\alpha = d\log(\lambda F(\lambda))/d\log\lambda$, calculated between 2.2 μm and 24 μm , is included in the catalogue for all source with at least two photometry points. In this paper we over-plot the positions of YSOs and protostars identified in this way, as well as the “red”, “red1” and “red2” sources (targets with a high MIPS band 1 to IRAC flux ratio, or those with rising spectral energy distributions [SEDs] detected in only MIPS1 or MIPS1 and IRAC band 4) and “PAH-em” and “star+dust” sources (targets with IRAC colours expected for PAH emission features; these may include background galaxies but also Class II sources). These latter sources are arguably also candidate protostars and/or outflow sources.

Note that both this paper and the *c2d* catalogue refer to all young stars as YSOs, regardless of whether they are Class 0/I protostars or Class II/III young stellar objects.

Sample SEDs, plotted between 1.2 μm and 24 μm , are presented for all source types by Evans et al. (2005). In Fig. 2 we plot a Spitzer colour-magnitude diagram for sources in Perseus West from the *c2d* catalogue. There are 66,535 sources in the catalogue, of which 4,352 are “red” sources and 610 YSOs. Clearly, most of the “red” sources, though red in the two IRAC bands, have a low 8.0 μm flux. In the most deeply embedded objects this may be caused by silicate absorption or the intrinsic shape of the SED (Class 0 sources have been shown to possess a [3.6]-[4.5]/[5.8]-[8.0] ratio that is >1 ; Jørgensen et al. 2006). Even so, the “red” sources may also include a modest population of background galaxies (Harvey et al. 2006; Jørgensen et al. 2006).

2.0.3 JCMT data

The H₂ emission in outflows is known to be produced in molecular shocks formed at the interface between a jet and the ambient medium. Thus, the H₂ knots and bow shocks imaged with WFCAM are expected to be closely related to the high-velocity molecular flow lobes often mapped

at (sub)millimetre wavelengths in rotational CO emission. In Figs.3 to 7 we therefore present CO J=3-2 observations of high-velocity molecular outflows in Perseus-West taken from the literature. These data were acquired at the James Clerk Maxwell Telescope (JCMT) and have been published and discussed in detail by Knee & Sandell (2000) and Hatchell, Fuller & Richer (2007a). The JCMT beam width measures 14'' at the 345.796 GHz frequency of the J=3 $\rightarrow$ 2 transition. Contour plots are presented in units of antenna temperature.

The maps cover only very limited regions in each star-forming cloud, though many of the flows discussed below have been observed. More extensive maps are being planned as part of the *Gould Belt* legacy survey at the JCMT (Ward-Thompson et al. 2007).

3 RESULTS & DISCUSSION

3.1 H₂ imaging of Perseus-West

The five WFCAM tiles used in this paper cover the western portion of Perseus, which includes NGC1333, L1455, L1448, and Barnard 1 (B1). WFCAM H₂ images of regions where outflows were detected are shown in Figs 3–8. The positions of protostars and YSOs identified in the Spitzer *c2d* survey are overplotted in each figure. Circles indicate sources that were given the “YSO” designation in the *c2d* catalogue, while triangles mark sources classified as “red”, “red2”, “PAH-em”, or “star+dust”. Jørgensen et al. (2007) have compared the *c2d* data with SCUBA observations at 850 μm . They present a catalogue of 49 “embedded” YSOs; all are MIPS detections that are either red in [3.6]-[4.5] and [8.0]-[24] colours or are located within 15'' of SCUBA peaks. They also include a few concentrated SCUBA cores with no MIPS counterparts. We label these sources as “YSO” with the numbers from Jørgensen et al.’s catalogue (their Table 3) and list those sources located in the Perseus-West region in Table 1.

The locations of Herbig-Haro (HH) objects found by Walawender, Bally & Reipurth (2005a) in their wide-field optical imaging, as well as HH objects that were previously known, are also marked on our figures. Note that these positions are often rather vague, representing the location of what is usually an extended object or diffuse feature.

In the online Appendix we present UKIRT-WFCAM/Spitzer-IRAC colour-composite images of each region, and discuss the outflows marked in Figs. 3–8 in some detail, noting in particular the association of H₂ emission-line features with HH objects, molecular CO outflows, dense cloud cores and the embedded source identified through Spitzer photometry. H₂ knot parameters (flow position angles, flow lengths, associated HH objects, etc.) are listed in Table A1, where we also give the embedded YSO that is likely to be driving each flow. Many of these knots or groups of knots are known HH objects or parts of outflows imaged in less extensive surveys. We therefore also list the names of associated HH objects and previous H₂ outflow names in Table A1.




Figure 3. Logarithmically-scaled H_2 $2.122\,\mu\text{m}$ image of B1 with the locations of HH objects marked with crosses, Spitzer-identified YSOs marked with circles and Spitzer-identified red sources marked with triangles. Jørgensen embedded YSOs are labelled. Contours of high-velocity CO $J=3-2$ emission (from Hatchell, Fuller & Richer 2007a) are over-plotted: integrated antenna temperature is plotted from 9 to $15\,\text{km s}^{-1}$ (dashed lines) and -5 to $4\,\text{km s}^{-1}$ (full lines). Contour levels are 1, 2, 4, 8 K km s^{-1} . H_2 features are numbered for reference (see Table A1). Outflows are indicated with arrows.




Figure 4. H_2 $2.122\,\mu\text{m}$ images of two adjacent regions in the B1 ridge. Contours of high-velocity CO 3-2 emission (from Hatchell, Fuller & Richer 2007a) are over-plotted from 9 to $15\,\text{km s}^{-1}$ (full lines) and from -5 to $4\,\text{km s}^{-1}$ (dashed lines). Contour levels are 0.5, 1, 2 K km s^{-1} (left panel) and 1, 1.5, 2 K km s^{-1} (right panel). See Fig.3 for further details.

3.2 On the merits of combining optical, infrared, and submillimetre observations of outflows

Overall, the combined WFCAM H_2 imaging, Spitzer imaging and photometry, and CO outflow observations provide

a clear and rather complete picture of dynamic activity in Perseus-West. Our WFCAM observations demonstrate that even shallow H_2 imaging is an effective tool for finding outflows from the youngest YSOs.

Of the 26 molecular outflows in our WFCAM tiles, 24




Figure 6. H_2 $2.122\,\mu\text{m}$ image of L1448. Contours of high-velocity CO 3-2 emission (from Hatchell, Fuller & Richer 2007a) are overplotted from 8 to $15\,\text{km s}^{-1}$ (full lines) and from -5 to $0\,\text{km s}^{-1}$ (dashed lines). Contour levels are $1, 2, 4, 8\,\text{K km s}^{-1}$. See Fig.3 for further details.

α may in part be due to excess emission from lines in some of the Spitzer bands (discussed below). However, when we plot flow length against Spitzer spectral index we find no correlation between these two parameters. This suggests that even the most deeply embedded H_2 jet source are old enough to produce parsec-scale flows. It seems likely that jet length, when derived from H_2 observations, is more sensitive to the environment (specifically the presence of ambient molecular gas that may be shock-excited into emission) than the “age” of the jet central engine.

Can we say anything about flow opening angles from our sample of jets in Perseus-West? High-resolution CO maps of outflows seem to indicate that Class 0 sources produce more collimated flows than their more evolved Class I counterparts (e.g. Lee et al. 2002; Arce & Sargent 2006). H_2 emission features (bow shocks) often envelope the outer edges of CO flow lobes, so one might expect to see a similar trend in H_2 data. However, shock-excited H_2 emission has a very short cooling time (of the order of a few years), so H_2 features are often short-lived and confined to a small percentage of the flow surface area. Moreover, close to the central source H_2 emission features tend to be associated with the central, on-axis jet rather than the swept-out cavity walls. Defining flow opening angles from H_2 images is therefore rather dif-

ficult. Even so, in Table 1 we list the mean opening angle, θ , for well-defined flows with multiple, bright H_2 emission features. θ is measured from cones in each flow lobe that are centred on the outflow source and which include all H_2 line-emission features. A plot of θ against source spectral index α (not shown) reveals no correlation between these two parameters. At best, we can only say that most of the flows in Perseus-West appear to be well collimated, as is expected for jets from young, embedded (high- α) sources. The opening angles listed have a mean of 12.5° and a standard deviation of only 4.1° . The associated sources have a relatively broad range in α : -0.87 to 3.79 .

Many of the H_2 jets in Perseus appear to be curved or possibly even precessing. If both lobes curve in the same direction (H_2 features 2–4 in the B1 ridge and 76/77 in L1448 are good examples) the flow curvature may be due to the motion of the source through the ambient medium. In such cases the source would need a proper motion that is a sizable fraction (at least a few percent) of the jet velocity. Source tangential velocities of the order of $1\text{--}10\,\text{km s}^{-1}$ would be required. Alternatively, the *apparent* curvature may simply be due to excitation of H_2 along only one side of an oval cavity, perhaps because of enhanced ambient gas densities




Figure 7. H_2 2.122 μm image of NGC1333. Open squares mark SCUBA cores without c2d-identified YSOs; the other symbols are as in Fig. 3. High-velocity CO 3-2 contours (from Knee & Sandell 2000) are over-plotted, from 12 to 18 $km\ s^{-1}$ (full lines) and from -5 to 3 $km\ s^{-1}$ (dashed lines). Contour levels are 1.5, 3, 6, 12 $K\ km\ s^{-1}$.

or an ambient density gradient across the width of the flow lobe.

Precession or jet “meandering” has been identified in a number of flows (e.g. Lopez et al. 1995; Heathcote et al. 1996; Davis et al. 1997; Hodapp et al. 2005). Precession may result from the misalignment of binary orbital and circum-binary disk axes (Terquem et al. 1999; Bate et al. 2000), although the disk should be forced into alignment with the orbital plane in roughly one precession period, which should be about 20-times longer than the binary orbital period (Bate et al. 2000). Given this time limitation, jet precession is expected from only the most embedded and therefore potentially very youngest sources. Notably, YSOs 11, 14 and 38, the three precessing jet candidates in Perseus-West, all have high values of α .

The H_2 flows from YSOs 11, 14 and 38 appear to have precessed through roughly half a turn. The dynamical age

of each flow should thus be a fair estimate of the precession period, to within a factor of ~ 2 (Hodapp et al. 2005). For a canonical jet velocity of 100 $km\ s^{-1}$ (consistent with jet proper motions, e.g. Eisloffel & Mundt 1994), from the flow lengths listed in Table 1, flow ages of 7300 yrs, 1400 yrs and 2600 yrs are estimated. Hence, a very crude range for the precession period of these remarkable flows is 2800-14,600 yrs.

3.4 Photometric characteristics of outflow sources

Fig. 9 demonstrates how the outflow sources differ from the bulk of the c2d-catalogue YSOs in Perseus. The outflow sources, plotted with black filled bars, are generally much redder and have a steeper near-IR/mid-IR SED than the general YSO population. The flow sources identified in this paper have a mean spectral index (α) of 1.39 ± 1.14 ; the c2d




Figure 8. H_2 2.122 μm image of the region to the south of NGC1333. Labelling and symbols are as in Fig.3.

YSOs, which are predominantly Class II T Tauri stars, have a mean α of -0.32 ± 0.89 . (In the c2d catalogue, in regions with $A_v < 5$ where sources are mostly background main sequence stars, the average value of α is -2.8 : Evans et al. 2005.)

The outflow sources also occupy a relatively distinct region in colour-colour space (Fig. 10). They possess a high $[4.5]-[24]$ IRAC/MIPS colour, as predicted for young sources with massive accretion disks (Whitney et al. 2004; Allen et al. 2004). However, they also exhibit positive $[3.6]-[4.5]$ colours, unlike the bulk of the c2d YSOs, where the near-zero $[3.6]-[4.5]$ colours are due to the intrinsic shape of the YSO SED. In Fig. 10, 80% of the outflow sources that have measured $3.6 \mu m$ and $4.5 \mu m$ magnitudes have a $[3.6]-[4.5]$ colour greater than 1. This is compared to only 11% of the total c2d catalogue of YSOs in Perseus. Of the YSOs with a $[3.6]-[4.5]$ colour greater than 1.0, 18% have outflows; only 0.4% of YSOs with $[3.6]-[4.5] < 1.0$ drive out-

flows. All factors point to the association of outflows traced in H_2 with the youngest Class 0/I protostars.

In Fig. 11 we plot α against $[3.6]-[4.5]$ colour for the entire Perseus c2d catalogue. Sources with values of α derived from just the two Spitzer bands plotted on the x-axis (i.e. those detected in only bands 1 and 2) lie along a narrow, diagonal strip. Sources detected in multiple Spitzer and/or 2MASS bands exhibit some scatter about this strip, since α is derived from photometry points that are modified by various absorption features (ice, silicate dust, etc.) and emission lines (H_2 , [FeII], [NeII], etc.), as well as the intrinsic shape of the YSO SED (i.e. that of an embedded source with a disk and probably also a disk-hole). The Spitzer YSOs tend towards higher values of α and redder $[3.6]-[4.5]$ colours, though they are largely indistinguishable from the other sources (field stars, background galaxies, etc.) in this plot. However, the outflow sources stand out as having excess emission at $4.5 \mu m$ and thus red $[3.6]-[4.5]$ colours. This is partly due to the inclusion of emission lines associated




Figure 9. Histograms of IRAC [3.6]-[4.5] colour and source spectral index, α , for YSOs in Perseus. Open bars represent all c2d identified YSOs; filled bars are sources with identified outflows.




Figure 10. Spitzer colour-colour diagrams for c2d YSOs in Perseus (open circles). Filled circles represent sources with identified outflows.

with infall and outflow in their integrated $4.5 \mu\text{m}$ fluxes. The $4.5 \mu\text{m}$ Spitzer band covers the H_2 pure-rotational 0-0S(9), S(10) and S(11) lines at 4.69 , 4.41 and $4.18 \mu\text{m}$, as well as the CO $v=1-0$ band between 4.45 - $4.95 \mu\text{m}$ and the HI Br α line at $4.052 \mu\text{m}$, both of which may be excited in accretion flows.

Some of the sources identified in the c2d catalogue as YSOs, based on their Spitzer colours and/or spectral index, will in fact be molecular hydrogen emission-line knots. Shock models developed by Smith & Rosen (2005) predict [3.6]-[4.5] colours in excess of ~ 0.5 , as well as positive values for α (when derived from all four IRAC bands). These predictions are supported by recent Spitzer observations of HH 46/47, where the jet and bow shocks are detected in all four IRAC channels (Velusamy, Langer & Marsh 2007) – note that there are pure-rotation H_2 emission lines in all of the IRAC bands.

In Fig. 11 we plot the [3.6]-[4.5] colours and corresponding spectral indices for six shock models. The arrows indicate

the effect of 5 magnitudes of K-band extinction (reddening moves the points to the upper-right in the figure). The arrow labelled “Planar” represents a slab of molecular gas at a temperature of 2000 K with a density of 10^5 cm^{-3} . Arrows labelled C80, C50, C40 and C20 are C-type bow shocks: all have the same shape and post-shock density (10^5 cm^{-3}), differing only in shock velocity (80 km s^{-1} , 50 km s^{-1} , 40 km s^{-1} and 20 km s^{-1} respectively). Decreasing shock velocity pushes the C-bow data points toward the upper-right in Fig. 11. Higher values of α and redder [3.6]-[4.5] colours are produced by a relative increase in emission from the lower-energy pure-rotational H_2 lines (0-0S(4) and S(5) in IRAC band 4 and 0-0S(6) and S(7) in band 3) over the higher energy 0-0 lines (S(9)-S(11) in band 2 and S(13) and above in band 1) and ro-vibrational lines (1-0O(5) and O(7) in band 1) at shorter wavelengths. The fastest C-type bow, C80, has a dissociative cap and is much like the J-type bow shock. It is therefore not too surprising that it coincides almost pre-




Figure 11. Plot of spectral index (α) against $[3.6]-[4.5]$ colour. Grey dots represent the entire c2d Perseus catalogue, red open circles represent c2d identified YSOs, and black filled circles represent outflow sources. Green circles mark c2d YSOs that may actually be misidentified H₂ knots. The arrows show the predicted positions for H₂ emission from a Planar shock, a J-type bow shock and four C-type bow shocks (Smith & Rosen 2005). The arrows reflect an increase in K-band extinction of 5 magnitudes.

cisely with the 50 km s^{-1} J-type bow model arrow, J50, in Fig. 11.

The shock model data points overlap quite significantly with the outflow sources in Fig. 11. The c2d “YSOs” that are most likely to be H₂ knots, features 7, 12, 13, 16, 38, 49 and 72 (the green circles in Fig. 11) have relatively low values of α and therefore appear to be associated with either fast C-type or J-type bow shocks. The low spectral indices measured for the observed knots could be caused by CO bandhead emission in IRAC band 2 or the inclusion of $24 \mu\text{m}$ data when calculating α for the observed knots. We do not include this data point in the model points, since there are no H₂ lines in the MIPS $24 \mu\text{m}$ bandpass. However, weak forbidden emission from [FeII] at $24.51 \mu\text{m}$ and $25.98 \mu\text{m}$ (like that seen in HH 46/47; Velusamy, Langer & Marsh 2007) could result in the “weak” detection of an outflow feature in this band in the c2d catalogue, and hence a lower value of α based on a fit to IRAC and MIPS data. CO emission is unlikely to contribute to observed outflow photometry data since very high densities, in excess of 10^6 cm^{-3} , are required to produce appreciable flux from this molecule. However, under such circumstances (seen perhaps in high-mass star forming regions) CO could account for as much as 10% of the observed emission in IRAC band 2.

3.5 Outflows and the ambient medium

The number of outflows identified here matches (to within 20%) the number of protostellar cores in L1448, L1455 and B1 (Hatchell et al. 2007b), though it is lower by a factor of 2 in NGC1333. In this densely-populated region this discrepancy is partly due to H₂ features being identified as part of a single flow, when in fact they are associated with multi-

ple flows. Even so, the association of protostellar cores with molecular (H₂/CO) outflows is clear.

It is also worth noting that the more massive clouds produce more outflows in Perseus-West. In B1, NGC1333, L1448 and L1455 we identify (at least) 8, 11, 4 and 5 outflows, respectively. From near-IR extinction mapping of Perseus, maps which cover an A_V range of about 3-7, Kirk et al. (2006) identify large scale structures which they refer to as “supercores”. They stress that the bulk of the cloud material is contained in these extensive, moderate-density regions, rather than in the high-extinction cores not traced by the extinction map. Kirk et al. measure masses of $441 M_\odot$, $973 M_\odot$, $174 M_\odot$ and $240 M_\odot$ for the B1, NGC1333, L1448 and L1455 supercores, respectively. Thus, in Perseus-West there is one molecular outflow for every 44-88 $M_\odot$ of ambient material. Roughly 50 generations of accretion/outflow activity would be needed to convert all of this molecular material into solar-mass stars.

In some regions many of the jets appear to be roughly parallel, though this common angle is not obviously related to the large-scale cloud structure. L1448 is perhaps the best example, where the position angles of the four outflows identified in Fig. 6 are between 147° and 155° . Kirk et al. (2006) fit 2-dimensional Gaussians to their supercores and, from their Fig. 5, we measure a supercore position angle of 120° for L1448. In L1448, the outflows thus appear to flow along the length of the supercore. However, the two parallel flows in the B1-ridge region (Fig. 4; PAs of 124° and 130°) are almost perpendicular to the elongated B1-ridge supercore (PA $\sim 50^\circ$). The flows in B1, NGC1333 and L1455 are more randomly orientated, although at least four of the larger flows in NGC1333 are aligned roughly north-south, parallel with the associated supercore (PA $\sim 10^\circ$). Even so, we conclude that in general the outflows are randomly orientated and that the large-scale cloud structure, as traced by the extinction mapping of Kirk et al. (2006), has no obvious influence on flow orientations.

3.6 Momentum injection and cloud support

Molecular outflows from young stars inject turbulent energy and momentum into the surrounding medium. Over time these flows probably provide turbulent support against cloud collapse and may even hinder the star formation process. In low-mass star forming regions usually only a dozen or so outflows are observed at any one time (NGC1333 is a good example). Here we investigate whether these outflows provide sufficient kinetic energy to account for the observed turbulence in the ambient medium.

As was noted by Walawender, Bally & Reipurth (2005a), one can estimate the turbulent momentum and energy in a star-forming region from the cloud mass, M_c , and velocity dispersion, δV_{tur} , measured from (sub)millimetre molecular lines:

- For Perseus-West we again use cloud masses from Kirk et al. (2006). The masses of their supercores (estimated from 2MASS extinction mapping for $A_V > 3$) are listed in Sect. 4.4. Although a sizable fraction of the overall cloud mass is likely to be in diffuse, lower- A_V regions, outflows are unlikely to impart much of their momentum into these extended regions. Also, masses derived from an extinction

map are preferred over masses estimated from molecular line observations, since the latter may suffer from depletion and/or opacity effects. And like submillimetre dust continuum maps, optically thin molecular lines will only trace the denser core regions ($A_V > 5$).

- A turbulent velocity is difficult to generalise for each region. The widths of molecular lines will result from turbulent and thermal motions in the molecular gas, as well as macroscopic motions if many independent cores are viewed along the same line of sight. In nearby, low-mass star forming regions the latter should not be a major problem. However, due to opacity and depletion effects, different molecular lines will trace different components of the ambient gas. Kirk, Johnstone & Tafalla (2007) find that in Perseus N₂H⁺ (1-0) lines are broadened almost exclusively by thermal motions in dense cores; C¹⁸O (2-1) is more sensitive to lower density material in the envelopes of dense cores (the C¹⁸O transition has a critical density of 10^3 cm^{-3} and freezes out onto dust grains at densities above 10^5 cm^{-3}). C¹⁸O is therefore a better tracer of the non-thermal motions in the interstellar – and intercore – environment pervaded by molecular outflows. Across Perseus C¹⁸O profile widths vary by only a few tenths of a km s^{-1} . From Kirk et al. (2007) we therefore adopt a mean value of 0.6 km s^{-1} for δV_{tur} for all regions considered.

In Table 2 we list the turbulent momentum $P_{\text{tur}} \sim M_c \delta V_{\text{tur}}$ and turbulent energy $E_{\text{tur}} \sim M_c \delta V_{\text{tur}}^2$ in each of the clouds observed in Perseus.

The momentum and energy of an outflow from a young star is hard to measure from infrared observations. Therefore, to estimate the combined momentum and energy, P_{jets} and E_{jets} , of the outflows from the embedded YSOs in each supercore in Perseus, we use the canonical values for the momentum flux for Class 0 and Class I low mass protostellar outflows measured by Bontemps et al. (1996), and multiply these by the accretion lifetime of the embedded phase (Walawender, Bally & Reipurth 2005a; Davis et al. 2007). The momentum and energy flux of an outflow is thought to decrease with time, in line with decreasing accretion rates (e.g. Smith 2000). We therefore ignore the Class II/III outflow phase. If the momentum flux for the Class 0 and I phases is $\sim 6 \times 10^{-5} \text{ M}_{\odot} \text{ km s}^{-1} \text{ yr}^{-1}$ and $\sim 4 \times 10^{-6} \text{ M}_{\odot} \text{ km s}^{-1} \text{ yr}^{-1}$ respectively, then for durations of 10^4 and 10^5 years, over its lifetime an outflow from a low-mass protostar should inject $\sim 1.0 \text{ M}_{\odot} \text{ km s}^{-1}$ of momentum into its surroundings. The outflow kinetic energy scales with the velocity; since most of the molecular material in outflows is probably entrained in the relatively slow-moving wings of bow shocks, then for a molecular flow velocity of $\sim 20 \text{ km s}^{-1}$ (e.g. Knee & Sandell 2000; Hatchell, Fuller & Richer 2007a), the kinetic energy supplied by a typical flow would be $\sim 4 \times 10^{37} \text{ J}$. In Table 2 we thus multiply these canonical values by the number of observed H₂ flows in each region to get P_{jets} and E_{jets} .

Although the combined energy of the outflows, E_{jets} , matches our estimates for the turbulent energy in each cloud, E_{tur} , the combined momentum, P_{jets} , falls short of P_{tur} by at least an order of magnitude in each region in Table 2. It has been argued that a large fraction of the jet momentum may be carried by a collimated, high-velocity jet (Walawender, Bally & Reipurth 2005a). However, the trans-

port of this momentum component to the surrounding cloud may be relatively inefficient, as the heavy jet punches its way through the ambient medium. The existence of parsec-scale jets and the high proper motions of even the most distant jet knots suggest that jets are relatively unhindered by their surroundings. Our assumption, that only the massive though low-velocity molecular flow component contributes to the turbulent motions in the ambient medium, is therefore probably still valid.

Instead, the turbulent momentum in the surrounding cloud may be built up via many generations of outflows. The lifetime of a Giant Molecular Cloud is 1-2 orders of magnitude longer than the duration of the Class 0/I phase, and Class 0, I and Class II YSOs clearly co-exist in many star-forming regions (e.g. Gutermuth et al. 2004; Kumar et al. 2007; Harvey et al. 2007; Gutermuth et al. 2008). Hence, over the lifetime of a GMC, many generations of outflows may contribute to the observed turbulent momentum in the surrounding cloud.

4 CONCLUSIONS

We examine near- and mid-IR images of a ~ 4 square degree region in Perseus-West and compare these to published catalogues of HH objects and moderately-extensive CO 3-2 maps. We find that in most cases molecular shock features are morphologically the same at $2.12 \mu\text{m}$ and $4.5 \mu\text{m}$, although the more embedded portions of some flows are revealed only in the longer-wavelength data. H₂ features are closely associated with high-velocity CO lobes, in support of bow shock entrainment scenarios. However, they often do not coincide with their optical HH counterparts, because of extinction effects or the discordant excitation requirements of each tracer.

We use IRAC and MIPS photometry to identify and characterise the outflow sources. In comparison to the full sample of YSOs in the c2d catalogue, the molecular outflow sources possess extreme values of near-to-mid-IR spectral index, α , and highly reddened [3.6]-[4.5] and [4.5]-[24] colours, as is befitting their extreme youth.

We find no correlation between H₂ flow length and α , nor between flow opening angle (as traced in H₂) and α . In general, the outflows are randomly orientated. However, we do find a correlation between the number of outflows and the number of protostellar cores in three of the four regions studied; in terms of total cloud mass, there is one molecular outflow for every 44-88 $\text{M}_{\odot}$ of ambient material. The outflows may also be an important source of turbulent energy in the interstellar medium.

ACKNOWLEDGEMENTS

Warm thanks are due to Nanda Kumar and Tom Megeath for advice on Spitzer data and YSO photometry, and to Jenny Hatchell and Lewis Knee for making their CO maps of Perseus available to us. We acknowledge the Cambridge Astronomical Survey Unit (CASU) for processing the near-IR data, and the WFCAM Science Archive in Edinburgh for making the WFCAM data discussed here available to us. The United Kingdom Infrared Telescope is operated by

the Joint Astronomy Centre on behalf of the U.K. Particle Physics and Astronomy Research Council. Some of the WFCAM data reported here were obtained in Directors Discretionary Time. This research made use of data products from the Spitzer Space Telescope Archive. These data products are provided by the services of the Infrared Science Archive operated by the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. We have also made extensive use of the SIMBAD database, operated at CDS, Strasbourg, France.

Table 1. Embedded YSOs and their molecular outflows in Perseus-West

Source ^a	H ₂ ^b features	CO ^b	HH ^b	Source Photometry and Spectral Index ^c						Length ^d (arcsec)	PA ^d (deg)	θ^d (deg)	Region
				[3.6]	[4.5]	[5.8]	[8.0]	[24]	α				
1	72,74	y	195	17.56	15.16	14.65	13.74	9.58	2.53	166	138	10	L1448
2	78-80,82-84	y	193,196,267	-	13.73	12.19	11.03	7.43	2.31	780	134	-	L1448
3	75-77	y	197	15.29	12.85	11.56	11.04	-	3.79	189	147	19	L1448
4	56-58	y	280,317,493,493	15.77	13.80	13.66	13.66	9.79	1.47	506	114	8	L1451
5	11-13,85?	n	n	11.76	11.59	11.52	11.43	9.96	-0.09	846	125	13	L1455
6	15-17,55	y	318	16.77	13.94	13.09	12.85	8.18	2.7	66	44	12	L1455
7	59	y	n	16.58	14.94	13.45	12.77	9.09	2.4	7	115	-	L1455
8	14,54	?	422	9.89	9.49	9.10	8.68	8.27	-0.19	37	179	8	L1455
9	66?	noOb	741,744?	15.28	16.18	15.28	14.92	11.92	0.68	-	-	-	NGC1333
10	n	n	n	14.61	13.57	12.86	12.56	11.84	0.16	-	-	-	NGC1333
11	20,24,46,71	noOb	15,338,339	13.24	10.87	10.59	9.31	-	3.16	461	140	-	NGC1333
12	n	n	n	18.54	17.68	17.02	16.47	12.81	1.71	-	-	-	NGC1333
13	61?	noOb	n	-	18.10	17.08	16.42	14.42	0.91	-	-	-	NGC1333
14	22,23,62	noOb	340,343	17.11	15.15	15.25	15.30	10.47	1.63	99	98	-	NGC1333
15	21,31,32,41,42,45	y	13,341,342	-	-	-	-	-	-	405	12	11	NGC1333
16	n	y	n	12.80	11.15	10.44	9.96	7.32	1.4	-	-	-	NGC1333
17	n	noOb	n	11.67	11.57	11.77	11.75	11.29	-0.74	-	-	-	NGC1333
18	26,27	noOb	n	-	19.03	18.30	-	13.29	2.18	127	98	10	NGC1333
19	38?	?	12?	-	-	-	-	-	-	-	-	-	NGC1333
20a ^e	35,60	y	7-11,753	8.90	11.67	7.53	19.77	-	0.37	114	132	17	NGC1333
20b ^e	30,37,38	y	12	-	-	-	-	-	-	235	172	5	NGC1333
20c ^e	19,28,34,43	n	14,344,350	-	-	-	-	-	-	987	10	4	NGC1333
21	n	n	n	17.13	16.15	16.01	15.96	11.65	1.55	-	-	-	NGC1333
22	29,33	y	n	-	18.92	18.62	-	12.57	2.63	89	34	9	NGC1333
23	39,40,44,48,65,67-70	y	5,347,761	-	14.12	13.80	13.89	9.09	1.9	412	159	12	NGC1333
24	36	y	6	-	14.80	14.92	15.33	9.50	2.18	27	48	58	NGC1333
25	n	n	n	17.41	13.58	15.31	15.51	12.44	0.72	-	-	-	NGC1333
26	n	n	n	19.32	16.69	16.52	-	12.67	1.8	-	-	-	NGC1333
28	n	noOb	n	16.34	15.15	14.34	13.83	11.73	1.06	-	-	-	NGC1333
29	n	noOb	n	12.22	12.23	11.98	11.42	10.59	-0.32	-	-	-	NGC1333
32	49,50	y	773	18.19	16.73	16.77	16.59	13.38	1.12	291	124	17	B1-ridge
33	2,3,4	y	782?	-	15.37	15.21	15.92	13.50	0.09	217	130	12	B1-ridge
35	n	y	n	18.28	16.32	15.52	14.80	11.16	2.17	-	-	-	B1
36	n	n	n	-	-	-	-	13.13	-	-	-	-	B1
37	10	y	n	14.46	12.03	11.46	11.21	8.31	1.9	13	90	53	B1
38	5-8	y	n	18.23	12.24	12.43	11.38	9.22	2.65	190	104	-	B1
39	9	y	n	14.75	13.22	12.78	12.59	10.76	0.67	45	62	18	B1
40	1	?	n	16.16	14.13	14.05	13.62	8.31	2.54	3	109	-	B1
L1448-IRS1	73,81	noOb	194	-	-	-	-	8.04	0.38	153	115	18	L1448
LkH α 327	63	noOb	432,356?	9.01	8.92	8.98	9.12	8.98	-0.87	142	151	6	B1

^aEmbedded YSOs in Perseus-West; list compiled by Jørgensen et al. (2007) from Spitzer and SCUBA observations (see their Table 3).^bH₂ features, molecular outflows, and HH objects associated with each YSO. See Table A1 for additional details.^cSource photometry obtained directly from the c2d catalogues.^dMean flow length, position angle and opening angle (average of two lobes).^eUnresolved YSOs associated with three separate, clearly defined flows (see the online Appendix for details).

Table 2. Turbulent momentum and energy in Perseus: clouds versus outflows

Region or supercore	P_{tur}^a ($M_{\odot}$ km s^{-1})	E_{tur}^a ($\times 10^{38}$ J)	N^b	P_{jets}^c ($M_{\odot}$ km s^{-1})	E_{jets}^c ($\times 10^{38}$ J)
B1	264	3.2	8	8	3.2
NGC1333	584	7.0	11	11	4.4
L1448	104	1.3	4	4	1.6
L1455	144	1.7	5	5	2.0

^aTurbulent momentum and energy associated with the molecular cloud.

^bNumber of H₂ flows in the region.

^cTotal momentum and energy associated with the “N” molecular outflows in each region.

REFERENCES

- Allen L.E. et al., 2004, ApJS, 154, 363
 Arce H.A., Sargent A.I., 2006, ApJ, 646, 1070.
 Aspin C., Sandell G., Russell A.P.G., A&AS, 106, 165.
 Bachiller R., Cernicharo J., Martin-Pintado J., Tafalla M., Lazareff B., 1990, A&A, 231, 174
 Bally J., Devine D., Hereld M., Rauscher B.J., 1993, ApJ, 418, L75
 Bally J., Devine D., Reipurth B., 1996, ApJ, 473, L49
 Bate M.R., Bonnell I.A., Clarke C.J., Lubow S.H., Ogilvie G.I., Pringle J.E., Tout C.A. 2000, MNRAS, 317, 773
 Bontemps S., Andre P., Tereby S., Cabrit S., 1996, A&A, 311, 858
 Casali M., et al., 2007, A&A 467, 777
 Černis K., 1990, Ap&SS, 166, 315
 Chrysostomou A., Hobson J., Smith M.D., Davis C.J., Berndsen A., 2000, MNRAS, 314, 229
 Dame, T. M., Hartmann D., Thaddeus P., ApJ, 2001, ApJ, 555, 12
 Davis C.J., Eisloffel J., Ray T.P., Jenness, T. 1997, A&A 324, 1013.
 Davis C.J., Kumar M.S.N., Sandell G., Froebrich D., Smith M.D., Currie M.J., 2007, MNRAS, 374, 29
 Davis C.J., Smith M.D., 1995, ApJ, 443, L41
 de Zeeuw P.T., Hoogerwerf R., de Bruijne J.H.J., Brown A.G.A., Blaauw A., 1999 AJ, 117, 354
 Dobashi K., Uehara H., Kandori R., Sakurai T., Kaiden M., Umemoto T., Sato F., 2005, PASJ, 57, 1
 Dye S., et al., 2006, MNRAS, 372, 1227
 Eisloffel J., 2000, A&A, 354, 236
 Eisloffel J., Mundt R., 1994, A&A, 284, 530
 Evans N.J. et al. 2003, PASP, 115, 965
 Evans N.J. et al. 2005, Delivery of Data from the c2d legacy Project: IRAC and MIPS (Pasedena: Spitzer science Center)
 Fazio G.G., et al., 2004 ApJS, 154, 10
 Gutermuth R.A., Megeath S.T., Muzerolle J., Allen L.E., Pipher J.L., Myers P.C., Fazio G., 2004, ApJS, 154, 374
 Gutermuth R.A., Myers P.C., Megeath S.T., Allen L.E., Pipher J.L., Muzerolle J., Porras A., Winston E., Fazio G., 2008, ApJ, in press.
 Harvey P.M., et al. 2006, ApJ, 644, 307.
 Harvey P.M., Merin B., Huard T.L., Rebull L.M., Chapman N., Evans N.J., Myers P.C., 2007, ApJ, 663, 1149.
 Hatchell J. et al., 2005, A&A, 440, 151
 Hatchell J., Fuller G.A., Richer J.S., 2007a, A&A, 472, 187
 Hatchell J., Fuller G.A., Richer J.S., Harries T., Ladd E.F., 2007b, A&A, 468, 1009
 Heathcote S., Morse J.A., hartigan P., Reipurth B., Schwartz R.D., Bally J., Stone J.M., 1996, AJ, 112, 3
 Hewett P.C., Warren S.J., Leggett S.K., Hodgkin S.T., 2006, MNRAS, 367, 454
 Hirano, N., Kamazaki, T., Mikami, H., Ohashi, N., & Umemoto, T. 1999, in Star Formation 1999, ed. T. Nakamoto (Nobeyama: NRO), 181
 Hodapp K.W., Bally J., Eisloffel J., Davis C.J., 2005, AJ, 129, 1580
 Hodapp K.W., Ladd E.F., 1995, ApJ, 453, 715
 Irwin M., et al., 2008, in prep.
 Jennings R.E., et al., 1987, MNRAS, 226, 461
 Jørgensen J.K. et al. 2006, ApJ, 645, 1246
 Jørgensen J.K. et al. 2007, ApJ, 656, 293
 Kirk H., Johnstone D., Di Francesco J., 2006, ApJ, 646, 1009
 Kirk H., Johnstone D., Tafalla M., 2007, ApJ, 668, 1042
 Knee L.B.G., Sandell G., 2000, A&A, 361, 671
 Kumar M.S.N, Davis C.J., Grave J.M.C., Ferreira B., Froebrich D., 2007, MNRAS, 374, 54.
 Lada C.J. et al., 2006, AJ, 131, 1574
 Ladd E.F., Lada E.A., Myers P.C., 1993, ApJ, 410, 168
 Lawrence A. et al., 2007, MNRAS, 379, 1599
 Lee C.-F., Mundy L.G., Stone J.M., Ostriker E.C., 2002, ApJ, 57, 294
 Lucas P.W. et al., 2008, MNRAS, in press.
 Luhman K.L., Rieke G.H., lada C.J., Lada E.A., 1998, ApJ, 508, 347
 Luhman K.L., et al., 2005, ApJ, 631, L69
 Lopez R., Raga A., Riera A., Anglada G., Estalella R., 1995, MNRAS, 274, L19
 Rieke G.H., et al. 2004, ApJSS, 154, 25
 Ridge N.A., Schnee S.L., Goodman A.A., Foster J.B. 2006, ApJ, 643, 932
 Rodriguez L.F., Anglada G., Curiel S., 1999, ApJSS, 125, 427
 Sargent A.I., 1979, ApJ, 233, 163.
 Smith M.D., 2000, Irish Astron. J., 27, 25
 Smith M.D., Rosen A., 2005, , ApJ, 233, 163.
 Takami M., et al. 2006, ApJ, 641, 357
 Terquem C., Eisloffel J., Papaloizou J.C.B., Nelson R.P., 1999, ApJ, 512, L131
 Tobin J., Looney L., Mundy M., Kwon W., Hamidouche M., 2007, ApJ, 659, 1404.
 Velusamy T., Langer W.D., Marsh K.A., 2007, ApJ, 668, L159
 Walawender J., Bally J., Kirk H., Johnstone D., 2005b, AJ, 130, 1795
 Walawender J., Bally J., Reipurth B., 2005a, AJ, 129, 2308
 Walawender J., Bally J., Reipurth B., Aspin C., 2004, AJ, 127, 2809
 Ward-Thompson D., et al., 2007, PASP, 119, 855
 Whitney B.A., et al., 2004, ApJS, 154, 315

APPENDIX A: DISCUSSION OF OUTFLOW ACTIVITY IN PERSEUS-WEST

In this section we present colour-composite images of each region constructed from WFCAM and IRAC observations, with $2.12\,\mu\text{m}$, $3.6\,\mu\text{m}$ and $4.5\,\mu\text{m}$ data occupying the blue, green and red channels, respectively (Figs A1–A5). These show how UKIRT/WFCAM and Spitzer/IRAC data complement each other in studies of outflows. H_2 features appear pink/purple, since line-emission is in almost all cases detected at $2.12\,\mu\text{m}$ and $4.5\,\mu\text{m}$. Higher-excitation H_2 knots will appear slightly bluer, since the WFCAM filter covers a rotation-vibrational line while the IRAC band 2 covers only pure-rotational transitions (discussed further in Sect. 4.3). The outflow sources, many of which are not visible in the near-IR, are in most cases detected in the IRAC bands. The most embedded sources appear red in the colour images, as do the most deeply embedded outflow features that are not detected at $2.12\,\mu\text{m}$.

These WFCAM/IRAC colour images are preferable to figures created from three of the four IRAC bands since one avoids using the less sensitive and relatively poor resolution IRAC bands 3 and 4 data: in colour composites made solely from IRAC data (e.g. $3.6/4.5/8.0\,\mu\text{m}$) faint, embedded sources may be undetected at the longer wavelength and thus will appear green, similar to H_2 line-emission features that are bright in IRAC band 2. The higher resolution WFCAM H_2 $2.122\,\mu\text{m}$ data also help to “sharpen” the outflow features, producing an effect similar to unsharp-masking.

Below we discuss the outflows in each region separately, relating the H_2 features detected at $2.12\,\mu\text{m}$ and/or $4.5\,\mu\text{m}$ with the high-velocity CO outflow lobes displayed in Figs. 3–7, the HH objects listed in the literature, and the protostars identified from Spitzer photometry (Evans et al. 2005; Jørgensen et al. 2007). Note that embedded YSOs from the list presented by Jørgensen et al. (2007) are referred to as YSO 1, YSO 2, etc., while sources from the c2d catalogue are identified by their RA and Dec, and are prefixed by “J”. Alternative source names are obtained from Table A1 of Hatchell et al. (2005) and references therein; NGC1333-IRAS source names are taken from Jennings et al. (1987).

A1 B1

We identify six outflows in B1. All six are driven by sources that are identified by c2d and all except LkHa327 are in the catalogue of Jørgensen et al. (2007). (There is potentially a seventh outflow that may have a source that is undetected by Spitzer between B1 and HH 429 - see below.)

The deeply embedded Class 0 source YSO 38 (SMM 1, B1-c) is associated with a well-defined bipolar CO outflow and the H_2 features 5 and 6-8 (Fig. 3). The H_2 data show that the outflow extends well beyond the bounds of the CO map. In our colour WFCAM/Spitzer image (Fig. A1), an ‘S-shaped’, precessing flow is evident; the flow is traced back to the source in the longer wavelength $4.5\,\mu\text{m}$ emission. YSO 38 clearly drives knot 5 and probably much of object 6-8 (see also Walawender et al. 2005b). Parts of the 6-8 complex could be powered by J033309.3+311012 (the YSO situated midway between features 6 and 7 in Fig. 3), though this source has a flat spectral index (~ 0.08) and only a moderately red $[3.6] - [4.5]$ colour (~ 0.80 ; see e.g.

Fig. 2). This source may instead be a bright outflow feature misidentified by c2d as a YSO candidate (discussed further in Sect. 4.3).

The overall morphology of the H_2 complex 6–8 also adds to the confusion. Its east-facing bow shape seems to suggest movement toward the east. 6–8 could conceivably be a bow shock in the counter-lobe of a flow that drives the bright, opposite-facing HH object HH 429 (H_2 knot 53; see Walawender et al. 2005b - Fig. 30). HH 429 is $17.5'$ due west of feature 53.

To the south of YSO 38, the nebulous, Class I source YSO 37 (Figs. 3 and A1) powers knot 10 to the west and a red-shifted CO lobe to the east. This east-west orientation supports the inclusion of HH 787, 785, 781 (and possibly HH 429) to the west and HH 789 to the east in another parsec-scale flow (Walawender et al. 2005b).

The blue-shifted CO emission to the southwest of YSO 37 is probably associated with a bipolar flow from YSO 35 (SMM3), although no H_2 emission is detected from this flow at $2.12\,\mu\text{m}$.

YSO 39 (B1-b, SMM2,3) comprises two Class 0 sources (B1-bN and B1-bS; Hirano et al. 1999). YSO 39 is associated with a bright, knotty H_2 flow 9 to the southwest, and a blue-shifted CO lobe to the northeast.

YSO 40 is a nebulous infrared source tentatively associated with blue-shifted CO emission and faint H_2 line emission (feature 50). The source has a relatively steep spectral index, as expected for an embedded outflow source ($\alpha = 2.54$, $[3.6] - [4.5] = 2.02$). It is possibly associated with the bright HH features HH 431,433 and the jet HH 790 (Walawender et al. 2005b, see their Fig. 32).

Finally, LkHa327, the brightest near-IR source in B1, drives a northwest-southeast outflow and excites HH 432 and H_2 feature 63. Walawender et al. (2005b) suggest that HH 430 and HH 788 trace the extended regions of the northern lobe, with HH 432 and 63 (their MH 7) tracing the southern lobe.

A2 B1 Ridge

The ridge of molecular gas running from B1 to L1455 contains two Jørgensen YSOs that are clearly driving molecular outflows (Figs. 4). YSO 33 (SMM 76, IRAS 03292+3039) is associated with H_2 features 2, 3 and 4, while YSO 32 (SMM 77, IRAS 03282+3035) powers H_2 features 49 (HH 773) and 50. Bow shock 49, first noted by Bally et al. (1993), is bright in H_2 $2.122\,\mu\text{m}$ (Fig. A2) but only marginally detected in the optical, as expected for embedded flows from Class 0 objects. The YSO 33 flow has no known HH counterparts.

51 and 52 are H_2 features located southwest along the B1 ridge (see Table A1 for details). The area has several Spitzer-identified YSOs and red sources, along with several HH objects which include HH 368-372, 427, 428, 769, and 792. However, none of the YSOs are in the catalogue of Jørgensen et al. (2007) (even though SCUBA data do exist for this region: Kirk et al. 2006). The HH objects and H_2 features do not have any clearly identifiable sources, although H_2 features 51, 52 and IRAS 03272+3013 do lie on the same axis.

Figure A1. SEE GIF UKIRT H₂ 2.122 μ m (blue) - Spitzer 3.6 μ m (green) - Spitzer 4.5 μ m (red) colour composite image of B1. H₂ features are numbered; embedded YSOs from Jørgensen et al. (2007) are labelled.

Figure A2. SEE GIF UKIRT H₂ 2.122 μ m (blue) - Spitzer 3.6 μ m (green) - Spitzer 4.5 μ m (red) colour composite image of B1 Ridge (west). H₂ features are numbered; embedded YSOs from Jørgensen et al. (2007) are labelled.

A3 L1455 and L1451

In the L1451/L1455 region in Fig. 5 we identify five H₂ outflows. At least four of these are driven by embedded YSOs (Jørgensen et al. 2007).

The most prominent outflow emerging from the L1455 core includes HH 279, H₂ knots 11-13 and possibly HH 423 and H₂ feature 85 in a parsec-scale flow (see Table 1). Class I objects YSO 5 (SMM 39) and YSO 8 (RNO 15) are candidate sources, although neither is associated with high-velocity CO (Fig. 5). The c2d source J032719.0+301718 (coincident with HH 279) has a red [3.6]-[4.5] colour of 1.55, though it also possess a negative spectral index (-0.46) and therefore could again be a H₂ knot mis-identified as a YSO. YSO 8 may instead drive a north-south flow that includes H₂ knots 14, 54 and weak blue-shifted CO emission to the south.

Between YSOs 5 and 8 in the southeast corner of Fig. 5, YSO 6 (SMM 35, RNO15-FIR, IRAS 03245+3002) powers H₂ features 15-17, 55 and a precessing bipolar CO outflow (Davis et al. 1997). YSO 7 (SMM 36) drives a second, orthogonal CO flow that includes the faint H₂ feature 59. The blue lobe of this CO outflow may extend $\sim 1'$ northwest of knots 15 and 16, although no H₂ features are detected in this extension. In the colour-composite image in Fig. A3 YSOs 6 and 7 are obvious as the reddest sources in the region.

To the west of L1455, toward L1451, there is a parsec scale flow driven by YSO 4 (SMM 80). The flow is associated with H₂ features 56, 57 and 58 (HH 492, 493 and 317, respectively) to the west, and the long chain of optical knots that comprise HH 280 to the east (Walawender et al. 2004). Red-shifted CO extends towards this eastern shock feature.

A4 L1448

There are four outflows identified in Fig. 6. The reddened source YSO 1 (SMM 30, IRAS 03222+3034, L1448-IRS2) drives a spectacular bipolar flow associated with H₂ features 72 and 74 (Fig. A4; see also Eisloffel 2000). Additional bright line emission features are seen to the northwest of YSO 1, although these are probably associated with a second bipolar flow from L1448-IRS1 (RNO 13). L1448-IRS1 is not catalogued as a YSO by Jørgensen. It has a spectral index of 0.38. It is detected only in the Spitzer MIPS 1 band in the c2d catalogue and is classified as “star+dust”. The lack of a detection in the IRAC bands is probably due to saturation as IRS1 is quite bright (see Fig. A4). H₂ knot 81 and HH 194 occupy the counter-lobe of this flow.

The parallel outflows from YSO 2 and YSO 3 (SMM 28/L1448N and SMM 29/L1448C, respectively) have been studied extensively in the past (e.g. Bachiller et al. 1990; Davis & Smith 1995; Eisloffel 2000, and references therein). The flow from YSO 2 includes H₂ features 82 to the southeast and 78, 79, 80, 83 and 84 to the northwest.

These features increase the length of this northwestern flow lobe to 25.5' (2.2 pc). The YSO 3 flow is associated with H₂ object 77 in the south and the spectacular, curving H₂ flow 76 to the north. Feature 75 is an unusual group of knots to the northwest of 76: we tentatively associate these with the curving 76 outflow, though on morphological grounds one might regard 75 and 78 as bow shocks and counter-bow shocks in a bipolar flow centred midway between these two objects. However, no candidate source is found in this region.

Finally, we bring the reader's attention to a highly reddened source 32'' south of YSO 3. This nebulous source can be seen clearly in the colour image in Fig. A4. The source, J032539.6+304336, is classified by c2d as “cup-down”, i.e. it has been detected in only 3 IRAC bands with a magnitude at 4.5 μ m that is brighter than at 3.6 μ m and 5.8 μ m (possibly due to unresolved line emission). However, close examination of the Spitzer IRAC images reveals that the source is in fact detected in all four IRAC bands. From our own photometry, measured from IRAC BCD images using a 2'' aperture, we measure a [3.6]-[4.5] colour of 2.2 (typical for outflow sources) and a spectral index, α , of 2.33 (again typical for embedded Class 0/I outflow sources). Hence, this source could be contributing to the outflow activity in this region.

A5 NGC1333

The outflows in NGC1333 have been studied in considerable detail, though the WFCAM images presented here are the most extensive H₂ images to date and as a result reveal new features. In Fig. 7 and 8 we mark 11 outflows. Some of these have previously been noted by Hodapp & Ladd (1995). All 11 are associated with Jørgensen et al. (2007) YSO sources.

A5.1 The SVS-13 SMM core

There are three major outflows emanating from the general area of SVS 13 and YSO 20. YSO 20 is identified by Jørgensen et al. (2007) as the centre of the SCUBA submillimeter core rather than a YSO. This is because YSO 20 was identified using only SCUBA data and so no c2d YSO position is available. This submillimeter core contains 5 radio continuum sources which Rodriguez et al. (1999) identify as YSOs.

Rather than speculate on the source of each of the flows emerging from the SVS 13 core, in Table 1, we identify the sources according to their alignment with outflow features and/or CO flow lobes: YSO 20a is deemed to be the driver of the well-know bipolar CO outflow containing HH 7-11; YSO 20b is the driver of the north-south flow at PA $\sim 170^\circ$ which includes H₂ features 30 in the south, 37 in the north and possible feature 38 (HH 12); YSO 20c is then the driver of the lone south lobe at PA $\sim 10^\circ$. This latter chain of H₂

Figure A3. SEE GIF UKIRT H₂ 2.122 μ m (blue) - Spitzer 3.6 μ m (green) - Spitzer 4.5 μ m (red) colour composite image of L1455. H₂ features are numbered; embedded YSOs from Jørgensen et al. (2007) are labelled.

Figure A4. SEE GIF UKIRT H₂ 2.122 μ m (blue) - Spitzer 3.6 μ m (green) - Spitzer 4.5 μ m (red) colour composite image of L1448. H₂ features are numbered; embedded YSOs from Jørgensen et al. (2007) are labelled.

features extends at least 16.5' (1.4 pc) to the south and includes H₂/HH objects 34, 28, 43, HH 14 and 19. There is no obvious counter-lobe in our H₂ image, although HH 754/755 do lie north along an axis drawn through the southern lobe (Walawender et al. 2005b). There is also no clear evidence of an outflow in the CO data of Knee & Sandell (2000). Our colour-composite image in Fig. A5 reveals a bright, nebulous source roughly 1' south of SVS 13 that could be associated with the YSO 20c flow. The source is associated with a conical reflection nebula that opens out toward the south.

A5.2 IRAS 2A and 2B

YSOs 15 and 16 are better known as IRAS 2A (SMM 44) and IRAS 2B. YSO 15 is associated with a SCUBA core and VLA radio source (Rodriguez et al. 1999). In the colour image in Fig. A5 YSO 15 appears highly reddened and elongated along the identified flow direction. It is clearly associated with H₂ knots 31 and 32, and the bipolar CO flow evident in Fig. 7. However, the YSO 15 “core” may also be associated with a second, orthogonal flow, which we label 45. This bipolar flow can also be seen clearly in the CO data. The north-south flow from YSO 15 is a large outflow that influences HH 12 in the north and extends at least as far down as HH 13 and H₂ features 41/42 in the south. Its axis also lines up with knot 18 (HH 746) $\sim$ 25' to the south of YSO 15. The extent of this flow lobe could therefore be at least 2.2 pc.

There are no H₂ features that are obviously associated with YSO 16 (IRAS 2B).

A5.3 HH 12

HH 12 is a complex cluster of knots about 5' north of SVS 13 and HH 7-11. The bright infrared source (IRAS 6) associated with HH 12 sits atop a column of dense, molecular gas; the source is flagged as a YSO by Jørgensen et al. (2007) based on its association with a SCUBA core (SMM 45).

We label the H₂ features around HH 12 as 37/38. Feature 37 is a faint trail of emission leading from near SVS 13 to the head of HH 12; 38 is the complex group of bright knots that make up the head of HH 12. Part of HH 12 could be associated with the H₂ flow 31/32 from YSO 15/IRAS 2A; HH 12 is clearly aligned with the well-defined axis of this bipolar outflow. However, the spray of knots to the west of YSO 19, coupled with the overlapping blue-shifted and red-shifted CO observed towards this source, point to an outflow inclined towards the line of sight that is driven by a source within HH 12. Rodriguez et al. (1999) find 2 radio continuum sources that coincide with the YSO 19 core and there are three flat-spectrum Spitzer YSOs about 1' to the southwest of YSO 19 that could be contributing to the outflow activity, although one or more of them could again be an H₂ knot misidentified as a c2d YSO.

A5.4 Other outflows in NGC1333

Northeast of SVS 13, H₂ feature 36 (HH 6) and its associated CO outflow are driven by either YSO 23 or YSO 24 (IRAS 7, SMM 46). There is also an extensive flow perpendicular to 36 that is likewise driven by either YSO 23 or 24. This large flow runs from H₂ knot 44/HH 5 in the south to feature 68 in the north (see Fig. A5). Feature 25/HH 18 and HH 761 may also be part of this flow. In Tables 1 and A1 YSO 23 is identified as the source of the extensive northeast-southwest jet and YSO 24 as the source of HH 6, because YSO 24 is slightly better aligned with the central axis of the feature 36/HH 6 cluster. However, the reverse cannot be ruled out.

NGC1333 also contains two “S-shaped” outflows, driven by YSOs 11 and 14 (Figs.7 and 8). YSO 11 (SMM 49) drives H₂ features 20, 46 and possibly 71/HH 338 (8.1' to the northwest), and feature 21 and possibly even 24/HH 15 (7.3' to the southeast). YSO 14 drives H₂ knots 22/23/62 and HH 340/343 in another S-shaped flow (for a detailed discussion of this outflow see Hodapp et al. 2005). Both flows are remarkably similar to the bending jet from YSO 38 in B1 (Fig. A1).

3' southeast of SVS 13, YSO 22 (SMM 41) drives a well-defined bipolar CO outflow that includes H₂ features 33 and 29 (Fig. 7). To the southwest, this flow may be interacting with the flow from the SVS 13 core and to the northeast with both the HH 7-11 outflow and the southern lobe of the YSO 23 jet.

Finally, near the bottom of Fig. 7, YSO 18 (SMM 65) drives an east-west flow associated with H₂ bows 26 to the east and 28 to the west. Unfortunately the CO map of Knee & Sandell (2000) does not extend sufficiently southward to contain YSO 18 and its outflow, or indeed any of the other southern flows labelled in Fig. 8.

Figure A5. SEE GIF UKIRT H₂ 2.122 μm (blue) - Spitzer 3.6 μm (green) - Spitzer 4.5 μm (red) colour composite image of NGC1333. H₂ features are numbered; embedded YSOs from Jørgensen et al. (2007) are labelled.

This paper has been typeset from a $\text{\TeX}$ / $\text{\LaTeX}$ file prepared by the author.

Table A1. H₂ features in Perseus-West

H ₂ feature	RA ^a (J2000.0)	Dec ^a (J2000.0)	Source ^b candidate	Length ^c (arcsec)	PA ^d (deg)	HH ^e	Comment	Other ^f identifiers	Figures ^g
1	3:33:27.417	31:07:10.06	40	3	109			MH 12	3,A1
2	3:32:11.860	30:50:51.55	33	243	-49				4
3	3:32:26.889	30:48:14.22	33	191	127				4
4	3:32:38.992	30:44:16.92	33	425	140				4
5	3:33:33.539	31:08:40.17	38	214	104			MH 1	3,A1
6	3:33:11.802	31:09:58.03	38	92	-71			part of MH 2	3,A1
7	3:33:07.400	31:10:16.64	38	154	-71			part of MH 2	3,A1
8	3:33:07.575	31:09:49.25	38	165	-76			part of MH 2	3,A1
9	3:33:18.648	31:07:14.35	39	45	-118			MH 5	3,A1
10	3:33:15.769	31:07:54.69	37	13	-90			MH 4	3,A1
11	3:27:01.380	30:18:27.11	5	606	-60		same flow as 12?		5,A3
12	3:27:16.405	30:17:14.44	5	408	-55	279			5,A3
13	3:27:20.454	30:17:15.97	5	258	-47				5,A3
14	3:27:47.874	30:12:24.62	8	44	-12				5,A3
15	3:27:38.776	30:12:56.34	6	25	-136				5,A3
16	3:27:37.773	30:12:46.72	6	25	-136		same as 6		5,A3
17	3:27:42.044	30:13:40.50	6	63	33				5,A3
18	3:28:32.601	30:50:30.12	-	57	161	746?	98'' from HH		8
19	3:28:49.421	31:00:11.13	20c	987	-171	14?	32'' from HH		8
20	3:28:36.965	31:13:31.42	11	100	-50	339?			7,A5
21	3:28:50.455	31:10:09.99	15	286	-166	341?			8
22	3:28:39.820	31:05:46.92	14	79	-83	340B			8
23	3:28:53.872	31:05:25.10	14	119	99	343			8
24	3:28:59.008	31:08:00.70	11	435	139	15			8
25	3:29:25.710	31:07:26.32	-	26	134	18			8
26	3:29:10.978	31:11:44.34	18	135	97			HL 8	7,A5
27	3:28:51.779	31:12:23.14	18	118	-80			ASR 97	7,A5
28	3:28:58.846	31:12:19.84	20c	223	-166	344?	66'' from HH	ASR 98	7,A5
29	3:29:06.977	31:12:15.71	22	88	-147			HL 3,5	7,A5
30	3:29:05.543	31:12:51.03	20b	193	174			HL 4	7,A5
31	3:28:53.911	31:13:24.08	15	76	-164			ASR 49,71	7,A5
32	3:28:56.054	31:15:47.67	15	74	10			ASR 15,16	7,A5
33	3:29:14.449	31:14:44.03	22	90	35	347?		ASR 57	7,A5
34	3:29:02.431	31:15:01.86	20c	54	-169			ASR 20,21	7,A5
35	3:29:00.695	31:16:58.24	20a	112	-35	753			7,A5
36	3:29:12.865	31:18:44.88	24	27	48	6			7,A5
37	3:28:59.208	31:18:51.44	20b	182	-10				7,A5
38	3:28:57.779	31:20:19.78	20b	265	-10	12			7,A5
39	3:29:03.812	31:21:57.75	23	237	-22				7,A5
40	3:29:00.448	31:24:04.77	23	369	-21				7,A5
41	3:28:46.819	31:07:08.72	15	464	-166	13			8
42	3:28:44.259	31:07:34.42	15	449	-161	13?	50'' from HH		8
43	3:28:56.631	31:07:37.47	20c	504	-170				8
44	3:29:20.190	31:12:49.95	23	352	160	5			7,A5
45	3:28:59.548	31:14:22.58	15	53	106			HL 2	7,A5
46	3:28:42.589	31:12:35.24	11	90	128				7,A5
47	3:29:27.277	31:13:30.71	-	6	63				7
48	3:29:10.873	31:19:53.97	23	85	0				7,A5
49	3:31:31.013	30:44:11.37	32	216	126	773	HH faint in optical		4,A2
50	3:31:02.864	30:47:47.50	32	365	-59				4,A2
51	3:30:33.338	30:27:51.29	-	21	155	370			-
52	3:30:45.780	30:29:19.27	-	10	154	372			-
53	3:31:47.973	31:10:00.60	-	47	94	429			-
54	3:27:47.978	30:11:44.37	8	30	180	422			5,A3
55	3:27:44.810	30:14:15.03	6	107	44	318			5,A3
56	3:26:49.129	30:14:56.63	4	165	105	493			5,A3
57	3:26:56.873	30:14:14.70	4	263	106	317D			5,A3
58	3:27:12.974	30:11:59.29	4	506	114	317			5,A3
59	3:27:43.006	30:12:29.64	7	7	115				5,A3

Table A1 – *continued*

MHF	RA ^a (J2000.0)	Dec ^a (J2000.0)	Source ^b candidate	Length ^c (arcsec)	PA ^d (deg)	HH ^e	Comment	Other ^f identifiers	Figure ^g
60	3:29:08.163	31:15:29.47	SVS 13	116	119	7-11		HL 7	7,A5
61	3:28:42.985	31:17:45.12	13						7,A5
62	3:28:42.879	31:05:36.47	14	32	99	340A	same flow as 22		8
63	3:33:35.820	31:08:45.76	LkH α 327	142	151			MH 7	3
64	3:33:13.323	31:08:18.11	-	5	90			MH 3	3
65	3:29:15.352	31:17:03.67	23	101	158				7,A5
66	3:28:33.274	31:12:14.42	-	8	163	744B			7,A5
67	3:29:17.212	31:15:26.51	23	192	154	374?		HL 10,11	7,A5
68	3:28:57.640	31:25:32.45	23	472	-22				-
69	3:29:03.949	31:23:02.04	23	295	-17				7,A5
70	3:29:07.627	31:21:04.69	23	170	-14				7
71	3:28:12.876	31:19:43.71	11	486	-40	338			-
72	3:25:26.718	30:44:04.20	1	122	140				6,A4
73	3:25:17.600	30:45:49.10	L1448-IRS1	138	112				6,A4
74	3:25:14.925	30:46:54.59	1	209	-44	195			6,A4
75	3:25:27.120	30:45:50.57	3	232	-53				6,A4
76	3:25:38.140	30:44:36.80	3	94	-30	197			6,A4
77	3:25:41.056	30:42:00.64	3	145	167				6,A4
78	3:25:12.777	30:49:22.86	2	390	-52	196			6,A4
79	3:24:52.682	30:54:47.39	2	863	-47	193			6
80	3:25:25.696	30:46:55.18	2	168	-56				6,A4
81	3:24:58.056	30:47:42.66	L1448-IRS1	168	-61	194			-
82	3:25:36.831	30:45:14.86	2	34	140				6,A4
83	3:24:26.679	30:56:44.31	2	1128	-53				-
84	3:24:02.837	31:00:25.00	2	1530	-52	267			-
85	3:28:40.874	30:01:59.92	5	1087	130				-

^aPosition of the brightest knot within the H_2 feature or flow.

^bMost likely source. Numbers refer to Table 3 in Jørgensen et al. (2007). A dash means there is no obvious source candidate.

^cThe distance from the source to the most distant knot. If no known source, entire length of all knots.

^dPosition angle, measured east of north, of an axis drawn from the source to the feature.

^eAssociated HH object (if any is known).

^fPrevious names of H_2 features. ASR refers to sources from Aspin, Sandell, & Russell (1994); HL refers to sources from Hodapp & Ladd (1995); MH refers to sources from Walawender et al. (2005b)

^gFigures in which the feature appears.

This figure "davis-fga1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0803.3075v1>

This figure "davis-fga2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0803.3075v1>

This figure "davis-fga3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0803.3075v1>

This figure "davis-fga4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0803.3075v1>

This figure "davis-fga5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/0803.3075v1>